

Single Crystal Analyses of ϵ -O₂ High-Pressure Phase of Solid Oxygen

Yuichi AKAHAMA(0001226)^{a*}, Haruki KAWAMURA(000127)^a

Osamu SHIMOMURA^b, Yukio NODA^b

^aFaculty of Science, Himeji Institute of Technology,

^bSPring-8 and ^cFaculty of Science, Chiba University

1. Introduction

Pressure-induced metallization and molecular dissociation of oxygen, O₂, with molecular magnetism have attracted special interest because of the novel electronic and magnetic properties of the high-pressure phases[1]. Determination of the structural properties of the high-pressure phases is indispensable for understanding the electro-magnetic properties.

In this study, the high-pressure single-crystal analysis of β -O₂ and ϵ -O₂ of low-Z element has been carried out including setup of the diffractometer on the BL02B1.

2. Experimental

The rhombohedral β -O₂ ($R\bar{3}m$) and monoclinic ϵ -O₂ crystals were grown under high pressure and temperature conditions at 5.5 GPa and 300 K and at 20 GPa and 600 K, respectively, in a Merrill-Bassett type DAC[2] which had a conical window of 70° ($2\theta=35^\circ$) to incident and scattering sides. The DAC was placed on a Huber diffractometer and the monochromated beam with an energy of 30 KeV($\lambda=0.4133 \text{ \AA}$) was exposed to the sample through a 0.5 mm diameter pin-hole.

3. Results and Discussion

For the β -O₂ sample, the UB parameter and the lattice constants of the hexagonal cell, $a=2.806(2)$ and $c=10.28(2)$ A, were determined from a least-squares fitting of 13 reflections. The values are listed in table I together with previous results[3]. The c -axis was unfortunately vertical to the top-surface of the diamond anvil. The typical reflection profile is shown in Fig.1. The almost reflections splitted into two or more peaks. Intensity data could be measured for 39 symmetry-independent reflections which were reduced to 12 nonequivalent

reflections. Structure refinement is in progress.

Two polycrystal 8-O2 samples with several crystals were tested with the oscillation photographs taken by using a vacuum camera and it was found that the single crystal analysis is feasible.

Table I. Structure parameters for β -O₂ phase together with the previous results[3].

		This work	Schiferl[3]
Pressure	$P(\text{GPa})$	5.7	5.5
Temperature	$T(\text{K})$	297	299
Lattice constant	$a(\text{\AA})$	2.806(2)	2.846
	$c(\text{\AA})$	10.285(2)	10.224
Volume	$V(\text{\AA}^3)$	70.2(2)	71.76
Atomic parameter	z		0.0577(2)
Thermal parameter	$\beta_{11}=\beta_{22}=2\beta_{12}$		0.1647(7)
	$\beta_{13}=\beta_{31}$		0
	β_{33}		0.0055(3)
Internuclear distance	$S(\text{\AA})$		1.202(2)
R factor	R_w		0.050




Fig. 1 Typical reflection profile of β -O₂ phase.

[1]Y.Akahama et al., Phys. Rev. Lett **74**, 4690(1995).

[2]Y.Akahama et al., in *Proc. AIRAPT Int. Conf.* Kyoto, (1997)p.781.

[3] D. Schiferl et al., Acta Cryst. **B37**, 1329 (1981).